

Catch-Up Guidance for Children 4 Months through 6 Years of Age

Diphtheria-, Tetanus-, and Pertussis-Containing Vaccines: DTaP/DT¹

The table below provides guidance for children whose vaccinations have been delayed. Start with the child's age and information on previous doses (previous doses must be documented and must meet minimum age requirements and minimum intervals between doses). Use this table in conjunction with table 2 of the Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, found at www.cdc.gov/vaccines/schedules/hcp/child-adolescent.html.

IF current age is	AND # of previous doses of DTaP or DT is	AND	THEN	Next dose due
4 months through 11 months	Unknown or 0	→	Give Dose 1 (DTaP) today	Give Dose 2 (DTaP) at least 4 weeks after Dose 1
	1	It has been at least 4 weeks since Dose 1	Give Dose 2 (DTaP) today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
		It has not been at least 4 weeks since Dose 1	No dose today	Give Dose 2 (DTaP) at least 4 weeks after Dose 1
	2	It has been at least 4 weeks since Dose 2	Give Dose 3 (DTaP) today	Give Dose 4 (DTaP) at least 6 calendar months after Dose 3 and at 15 months of age or older ²
		It has not been at least 4 weeks since Dose 2	No dose today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
	1 through 3 years	Unknown or 0	→	Give Dose 1 (DTaP) today
1		It has been at least 4 weeks since Dose 1	Give Dose 2 (DTaP) today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
		It has not been 4 weeks since Dose 1	No dose today	Give Dose 2 (DTaP) at least 4 weeks after Dose 1
2		It has been at least 4 weeks since Dose 2	Give Dose 3 (DTaP) today	Give Dose 4 (DTaP) at least 6 calendar months after Dose 3
		It has not been 4 weeks since Dose 2	No dose today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
3		It has been at least 6 calendar months since Dose 3	If 12 through 14 months of age, no dose today ²	Give Dose 4 (DTaP) at 15 through 18 months of age
			If 15 months of age or older, give Dose 4 (DTaP) today	Give Dose 5 (DTaP) at least 6 months after Dose 4 and at 4 through 6 years of age
		It has not been 6 calendar months since Dose 3	No dose today	Give Dose 4 (DTaP) at least 6 months after Dose 3

¹Vaccine information: DTaP—Administer to children 6 weeks through 6 years of age without a contraindication or precaution to diphtheria, tetanus, or pertussis vaccine. DTaP products include Daptacel, Kinrix, Infanrix, Pediarix, Pentacel, and Quadracel. Use the correct product based on the approved age indications. DT—Administer to children 6 weeks through 6 years of age with a contraindication to pertussis vaccine.

²The fourth dose may be administered as early as age 12 months, provided at least 6 months have elapsed since the third dose.

Reference: Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger—United States, 2019. www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

Catch-Up Guidance for Children 4 Months through 6 Years of Age

Diphtheria-, Tetanus-, and Pertussis-Containing Vaccines: DTaP/DT¹

IF current age is	AND # of previous doses of DTaP or DT is ¹	AND	AND	THEN	Next dose due
4 through 6 years	Unknown or 0	→	→	Give Dose 1 (DTaP) today	Give Dose 2 (DTaP) at least 4 weeks after Dose 1
	1	It has been at least 4 weeks since Dose 1	→	Give Dose 2 (DTaP) today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
		It has not been 4 weeks since Dose 1	→	No dose today	Give Dose 2 (DTaP) at least 4 weeks after Dose 1
	2	It has been at least 4 weeks since Dose 2	→	Give Dose 3 (DTaP) today	Give Dose 4 (DTaP) at least 6 calendar months after Dose 3
		It has not been at least 4 weeks since Dose 2	→	No dose today	Give Dose 3 (DTaP) at least 4 weeks after Dose 2
	3	It has been at least 6 calendar months since Dose 3	→	Give Dose 4 (DTaP) today	Give Tdap at 11 to 12 years of age
		It has not been at least 6 calendar months since Dose 3	→	No dose today	Give Dose 4 (DTaP) at least 6 calendar months after Dose 3
	4	All doses were given prior to the 4 th birthday	It has not been at least 6 months since Dose 4	No dose today	Give Dose 5 (DTaP) at least 6 calendar months after Dose 4
			It has been at least 6 months since Dose 4	→	Give Dose 5 (DTaP) today
		At least one dose was given at/after the 4 th birthday	→	No dose today	Give Tdap at 11 to 12 years of age

¹Vaccine information:DTaP—Administer to children 6 weeks through 6 years of age without a contraindication or precaution to diphtheria, tetanus, or pertussis vaccine. DTaP products include Daptacel, Kinrix, Infanrix, Pediarix, Pentacel, and Quadracel. Use the correct product based on the approved age indications. DT—Administer to children 6 weeks through 6 years of age with a contraindication to pertussis vaccine.

Reference: Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger—United States, 2019.
www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf